

ON THE $PSL_2(\mathbb{F}_{19})$ -INVARIANT CUBIC SEVENFOLD

ATANAS ILIEV, XAVIER ROULLEAU

ABSTRACT. It has been proved by Adler that there exists a unique cubic hypersurface X^7 in $\mathbb{P}^8$ which is invariant under the action of the simple group $PSL_2(\mathbb{F}_{19})$. In the present note we study the intermediate Jacobian of X^7 and in particular we prove that the subjacent 85-dimensional torus is an Abelian variety. The symmetry group $G = PSL_2(\mathbb{F}_{19})$ defines uniquely a G -invariant abelian 9-fold $A(X^7)$, which we study in detail and describe its period lattice.

INTRODUCTION

Let $PSL_2(\mathbb{F}_q)$ be the projective special linear group of order 2 matrices over the finite field with q element $\mathbb{F}_q$. There exist exactly two non trivial irreducible complex representations $W_{\frac{q-1}{2}}, \bar{W}_{\frac{q-1}{2}}$ of $PSL_2(\mathbb{F}_q)$ on a space of dimension $\frac{q-1}{2}$, each one complex conjugated to the other, see [3]. In [1], Adler proved that for any q a power of a prime $p \geq 7$, with $p = 3 \bmod 8$, there exists (up to a constant multiple) a unique $PSL_2(\mathbb{F}_q)$ -invariant cubic form f_q of $W_{\frac{q-1}{2}}$. We call the corresponding unique $PSL_2(\mathbb{F}_q)$ -invariant cubic hypersurface

$$X^{\frac{q-5}{2}} = \{f_q = 0\} \hookrightarrow \mathbb{P}(W_{\frac{q-1}{2}})$$

the *Adler cubic* for q .

In the smallest non-trivial case, $q = 11 = 1 \cdot 8 + 3$, the the Adler cubic in $\mathbb{P}^4 = \mathbb{P}(W_{\frac{11-1}{2}})$ coincides with the Klein cubic threefold

$$X^3 = \{x_1^2 x_2 + x_2^2 x_3 + x_3^2 x_4 + x_4^2 x_5 + x_5^2 x_1 = 0\}.$$

This threefold has been introduced by Felix Klein as a higher dimensional analog of the well known Klein quartic curve, the unique curve in the projective plane with symmetry group $PSL_2(\mathbb{F}_7)$, see [14]. In [17], the second author has proven that there exists a unique Abelian fivefold $A(X^3)$ with a $PSL_2(\mathbb{F}_{11})$ -invariant principal polarization and he has explicitly described the period lattice of $A(X^3)$. In this case, $A(X^3)$ coincide with the Griffiths intermediate Jacobian $J_G(X^3)$ of the cubic threefold X^3 , with the principal polarization coming from the intersection of real 3-cycles on X^3 .

2000 *Mathematics Subject Classification.* 11G10, 14J50, 14J70.

Key words and phrases. Cubic sevenfold, automorphism group, Intermediate Jacobian.

In this paper we study the next case - the Adler cubic for $q = 19 = 2 \cdot 8 + 3$. By using the general descriptions from Theorem 4 of [1], one can find an equation of the Adler cubic X^7 :

$$f_{19} = x_1^2 x_6 + x_6^2 x_2 + x_2^2 x_7 + x_7^2 x_4 + x_4^2 x_5 + x_5^2 x_8 + x_8^2 x_9 + x_9^2 x_3 + x_3^2 x_1 \\ - 2(x_1 x_7 x_8 + x_2 x_3 x_5 + x_4 x_6 x_9).$$

In the first section, we study a similar invariant principally polarized Abelian ninefold $A(X^7)$ defined uniquely by the Adler cubic sevenfold, and compute the period lattice and the first Chern class of the polarization of $A(X^7)$.

In the second section, we show that on the 85-dimensional complex torus $J = (H^{5,2} \oplus H^{4,3})^* / H_7(F_{19}, \mathbb{Z})$ of the Griffiths intermediate Jacobian $J_G(X^7)$ of X^7 , one can introduce a structure of a $PSL_2(\mathbb{F}_{19})$ -invariant polarization which provides J with a structure of an Abelian variety.

In the third section, we study invariant properties of the Adler-Klein pencil of cubic sevenfolds, an analog of the Dwork pencil of quintics threefolds see e.g. [9]. Notice that from the point of view of variation of Hodge structure, the cubic sevenfolds can be considered as higher dimensional analogs of Calabi-Yau threefolds, see [7] and [13].

Acknowledgements. We thank Bert van Geemen for its comments on a preliminary version of this paper. The second author acknowledges the hospitality of the Seoul National University where part of this work was done. He is a member of the project Geometria Algebraica PTDC/MAT/099275/2008 and was supported by FCT grant SFRH/BPD/72719/2010.

1. THE INVARIANT ABELIAN 9-FOLD OF THE ADLER CUBIC 7-FOLD.

We begin by some notations. Let $e_1, \dots, e_9$ be a basis of a 9-dimensional vector space V . Let $\tau \in GL(V)$ be the order 19 automorphism defined by

$$\tau : e_j \rightarrow \xi^{j^2} e_j,$$

where $\xi = e^{2i\pi/19}$, $i^2 = -1$, and let be $\sigma \in GL(V)$ the order 9 automorphism defined by

$$\sigma : e_k \rightarrow e_{|6j|},$$

where for any integer t the notation $|t|$ means the unique integer $0 \leq u \leq 9$ such that $t = \pm u \bmod 19$. We denote by μ the order 2 automorphism given in the basis $e_1, \dots, e_9$ by the matrix:

$$\mu = \left(\frac{i}{\sqrt{19}} \left(\frac{kj}{19} \right) (\xi^{kj} - \xi^{-kj}) \right)_{1 \leq k, j \leq 9}$$

where $\left(\frac{kj}{19} \right)$ is the Legendre symbol. The group generated by τ, σ and μ is isomorphic to $PSL_2(\mathbb{F}_{19})$ and define a representation of $PSL_2(\mathbb{F}_{19})$, see [3].

Let us define

$$v_k = \tau^k(e_1 + \cdots + e_9) = \xi^k e_1 + \xi^{4k} e_2 + \xi^{9k} e_3 + \xi^{16k} e_4 + \xi^{6k} e_5 + \xi^{17k} e_6 + \xi^{11k} e_7 + \xi^{7k} e_8 + \xi^{5k} e_9$$

(thus $v_k = v_{k+19}$) and

$$w'_k = \frac{1}{1+2\nu}(v_k - 5v_{k+1} + 10v_{k+2} - 10v_{k+3} + 5v_{k+4} - v_{k+5}),$$

where

$$\nu = \sum_{k=1}^{k=9} \xi^{k^2} = \frac{-1 + i\sqrt{19}}{2}.$$

An endomorphism h of a torus A acts on the tangent space $T_{A,0}$ by its differential dh , which we call the *analytic representation* of h . For the ease of the notations, we will use the same letter for h and its analytic representation.

In the present section, we will prove the following Theorem:

Theorem 1. *There exist:*

(1) a 9-dimensional torus $A = V/\Lambda$, $T_{A,0} = V$ such that the elements τ , σ of $GL(V)$ are the analytic representations of automorphisms of A . The torus A is isomorphic to the Abelian variety E^9 , where E is the elliptic curve $\mathbb{C}/\mathbb{Z}[\nu]$, ($\nu = \frac{-1+i\sqrt{19}}{2}$).

(2) a unique principal polarization Θ on A which is invariant by the automorphisms σ, τ . The period lattice of A is then:

$$H_1(A, \mathbb{Z}) = \frac{\mathbb{Z}[\nu]}{1+2\nu} w'_0 + \frac{\mathbb{Z}[\nu]}{1+2\nu} w'_1 + \frac{\mathbb{Z}[\nu]}{1+2\nu} w'_2 + \frac{\mathbb{Z}[\nu]}{1+2\nu} w'_3 + \bigoplus_{k=4}^8 \mathbb{Z}[\nu] v_k,$$

and $c_1(\Theta) = \frac{2}{\sqrt{19}} \sum_{k=1}^{k=9} dx_k \wedge d\bar{x}_k$, where $x_1, \dots, x_9$ is the dual basis of the e_j 's.

In order to prove Theorem 1, we first suppose that such a torus $A = V/\Lambda$ exists and find the necessary condition for its existence. We then check immediately that these conditions are also sufficient, and that they give us the uniqueness of A .

Let $\Lambda \subset V$ be a lattice such that τ and σ of $GL(V)$ are analytic representations of automorphisms of the torus $A = V/\Lambda$. Let

$$\ell_k = \xi^k x_1 + \xi^{4k} x_2 + \xi^{9k} x_3 + \xi^{16k} x_4 + \xi^{6k} x_5 + \xi^{17k} x_6 + \xi^{11k} x_7 + \xi^{7k} x_8 + \xi^{5k} x_9.$$

Let q be the endomorphism $q = \sum_{j=0}^{j=8} \sigma^j$. For $z = \sum_{j=1}^{j=9} x_j e_j$, we have $q(z) = \ell_0(z) v_0$, thus the image of q is an elliptic curve $\mathbb{E}$ contained in A .

The restriction of $q \circ \tau : A \rightarrow \mathbb{E}$ to $\mathbb{E} \hookrightarrow A$ is the multiplication by $\nu = \ell_0(\tau v_0)$. Thus the endomorphism group $\text{End } \mathbb{E}$ of $\mathbb{E}$ contains the ring $\mathbb{Z}[\nu]$; since this is a maximal order, we have $\text{End } \mathbb{E} = \mathbb{Z}[\nu]$. We remark that the ring $\mathbb{Z}[\nu]$ is one of the 9 rings of integers of quadratic fields that are

Principal Ideal Domains, see [15]. Since $\mathbb{Z}[\nu]$ is a PID and $H_1(A, \mathbb{Z}) \cap \mathbb{C}v_0$ is a rank one $\mathbb{Z}[\nu]$ -module, there exists a constant $c \in \mathbb{C}^*$ such that

$$H_1(A, \mathbb{Z}) \cap \mathbb{C}v_0 = \mathbb{Z}[\nu]cv_0.$$

Up to normalization of the e_j 's, we can suppose that $c = 1$.

Let $\Lambda_0 \subset V$ be the $\mathbb{Z}$ -module generated by the v_k , $k \in \mathbb{Z}/19\mathbb{Z}$. The group Λ_0 is stable under the action of τ and $\Lambda_0 \subset H_1(A, \mathbb{Z}) = \Lambda$.

Lemma 2. *The $\mathbb{Z}$ -module $\Lambda_0 \subset H_1(A, \mathbb{Z})$ is equal to the lattice:*

$$R_0 = \sum_{k=0}^{k=8} \mathbb{Z}[\nu]v_0.$$

Proof. We have $\nu v_0 = \sum_{k=1}^{k=9} v_k$ hence νv_0 is an element of Λ_0 . This implies that the vectors $\nu v_k = \tau^k \nu v_0$ are elements of Λ_0 for all k , hence: $R_0 \subset \Lambda_0$. Conversely, we have:

$$v_9 = v_0 + (1+\nu)v_1 - 2v_2 + (1-\nu)v_3 + (3+\nu)v_4 + (-2+\nu)v_5 - (2+\nu)v_6 + 2v_7 + \nu v_8$$

and similar formulas for $v_{10}, \dots, v_{17}$. This proves that the lattice R_0 contains the vectors $v_k = \tau^k v_0$ generating Λ_0 , thus: $R_0 = \Lambda_0$. $\square$

Now we construct a lattice that contains $H_1(A, \mathbb{Z})$. Let be $k \in \mathbb{Z}/19\mathbb{Z}$. The image of $z \in V$ by the endomorphism $q \circ \tau^k : V \rightarrow V$ is

$$q \circ \tau^k(z) = \ell_k(z)v_0.$$

Let be $\lambda \in H_1(A, \mathbb{Z})$. Since $H_1(A, \mathbb{Z}) \cap \mathbb{C}v_0 = \mathbb{Z}[\nu]v_0$, the scalar $\ell_k(\lambda)$ is an element of $\mathbb{Z}[\nu]$. Let

$$\Lambda_8 = \{z \in V \mid \ell_k(z) \in \mathbb{Z}[\nu], 0 \leq k \leq 8\}.$$

Lemma 3. *The $\mathbb{Z}$ -module $\Lambda_8 \supset H_1(A, \mathbb{Z})$ is the lattice:*

$$\sum_{k=0}^{k=7} \frac{\mathbb{Z}[\nu]}{1+2\nu} (v_k - v_{k+1}) + \mathbb{Z}[\nu]v_0.$$

Moreover τ stabilizes Λ_8 .

Proof. Let $\ell_0^*, \dots, \ell_8^*$ be the basis dual to $\ell_0, \dots, \ell_8$. By definition, the $\mathbb{Z}[\nu]$ -module Λ_8 is $\oplus_{i=0}^{i=8} \mathbb{Z}[\nu]\ell_i^*$. By expressing the ℓ_i^* in the basis $v_0, \dots, v_8$ we obtain the lattice Λ_8 . Using the formula:

$$\begin{aligned} v_9 - v_8 &= v_0 + (1+\nu)v_1 - 2v_2 + (1-\nu)v_3 + (3+\nu)v_4 \\ &\quad + (\nu-2)v_5 - (2+\nu)v_6 + 2v_7 + (\nu-1)v_8, \end{aligned}$$

one can check that $v_9 - v_8$ is a $\mathbb{Z}[\nu]$ -linear combination of $(2\nu+1)v_0$ and the $v_k - v_{k+1}$ for $k = 0, \dots, 7$. Therefore $\tau(\frac{1}{2\nu+1}(v_8 - v_7)) = \frac{1}{2\nu+1}(v_9 - v_8)$ is in Λ_8 and Λ_8 is stable by τ . $\square$

We denote by $\phi : \Lambda_8 \rightarrow \Lambda_8/\Lambda_0$ the quotient map. The ring $\mathbb{Z}[\nu]/(1+2\nu)$ is the finite field with 19 elements. The quotient Λ_8/Λ_0 is a $\mathbb{Z}[\nu]/(1+2\nu)$ -vector space with basis $t_1, \dots, t_8$ such that $t_i = \frac{1}{1+2\nu}(v_{i-1} - v_i) + \Lambda_0$.

Let R be a lattice such that $\Lambda_0 \subset R \subset \Lambda_8$. The group $\phi(R)$ is a sub-vector space of Λ_8/Λ_0 and:

$$\phi^{-1}\phi(R) = R + \Lambda_0 = R.$$

The set of such lattices R correspond bijectively to the set of sub-vector spaces of Λ_8/Λ_0 .

Because the automorphism τ preserves Λ_0 , it induces an automorphism $\hat{\tau}$ on the quotient Λ_8/Λ_0 such that $\phi \circ \tau = \hat{\tau} \circ \phi$. As τ stabilizes $H_1(A, \mathbb{Z})$, the vector space $\phi(H_1(A, \mathbb{Z}))$ is stable by $\hat{\tau}$. Let

$$w_{8-k} = (-1)^k \sum_{j=0}^{j=k} \binom{k}{j} (-1)^j t_{j+1}, \quad k = 0 \dots 7.$$

Then $w_{i-1} = \hat{\tau}w_i - w_i$ for $i > 1$ and $\hat{\tau}w_1 = w_1$. The matrix of $\hat{\tau}$ in the basis $w_1, \dots, w_8$ is the size 8×8 matrix:

$$\begin{pmatrix} 1 & 1 & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & \ddots & \ddots & 1 \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}$$

The sub-spaces stable by $\hat{\tau}$ are the spaces W_j , $1 \leq j \leq 8$ generated by $w_1, \dots, w_j$ and $W_0 = \{0\}$. Let Λ_j be the lattice $\phi^{-1}W_j$. It is easy to check that, as a lattice in $\mathbb{C}^9$, the Λ_j are all isomorphic to $\mathbb{Z}[\nu]^9$. Since $\Lambda = H_1(A, \mathbb{Z})$ is stable by τ , we have proved that :

Lemma 4. *The torus $A = V/\Lambda$ exists, it is an Abelian variety isomorphic to E^9 . There exists $j \in \{0, 1, \dots, 8\}$ such that $\Lambda = \Lambda_j$.*

Let $w'_0, \dots, w'_3$ be the vectors defined by:

$$w'_0 = \frac{1}{1+2\nu}(v_0 - 5v_1 + 10v_2 - 10v_3 + 5v_4 - v_5) = \frac{1}{1+2\nu}(1 - \tau)^5 v_0$$

and $w'_k = \tau^k w'_0$ (we have $\phi(w'_0) = w_4$). Let us suppose that $A = V/\Lambda$ has moreover a principal polarization Θ that is invariant under the action of τ and σ . Then:

Lemma 5. *The lattice $H_1(A, \mathbb{Z})$ is equal to Λ_4 , and*

$$\Lambda_4 = \frac{\mathbb{Z}[\nu]}{1+2\nu}w'_0 + \frac{\mathbb{Z}[\nu]}{1+2\nu}w'_1 + \frac{\mathbb{Z}[\nu]}{1+2\nu}w'_2 + \frac{\mathbb{Z}[\nu]}{1+2\nu}w'_3 + \bigoplus_{k=4}^8 \mathbb{Z}[\nu]v_k.$$

The Hermitian matrix associated to Θ is equal to $\frac{2}{\sqrt{19}}I_9$ in the basis $e_1, \dots, e_9$ and $c_1(\Theta) = \frac{i}{\sqrt{19}} \sum_{k=1}^{k=9} dx_k \wedge d\bar{x}_k$.

Proof. Let H be the matrix of the Hermitian form of the polarization Θ in the basis $e_1, \dots, e_9$. Since τ preserves the polarization Θ , this implies that:

$${}^t\tau H \bar{\tau} = H$$

where $\bar{\tau}$ is the matrix whose coefficients are conjugated of those of τ . The only Hermitian matrices that verify this equality are the diagonal matrices. By the same reasoning with σ instead of τ , we obtain that these diagonal coefficients are equal, and:

$$H = a \frac{2}{\sqrt{19}} I_9,$$

where a is a positive real (H is a positive definite Hermitian form). As H is a polarization, the alternating form $c_1(\Theta) = \Im m(H)$ takes integer values on $H_1(A, \mathbb{Z})$, hence $\Im m({}^t v_2 H \bar{v}_1) = a$ is an integer.

Let $c_1(\Theta) = \Im m(H) = i \frac{a}{\sqrt{19}} \sum dx_k \wedge d\bar{x}_k$ be the alternating form of the principal polarization Θ . Let $\lambda_1, \dots, \lambda_{18}$ be a basis of a lattice Λ' . By definition, the square of the Pfaffian $Pf_{\Theta}(\Lambda')$ of Λ' is the determinant of the matrix

$$M_{\Lambda'} = (c_1(\Theta)(\lambda_j, \lambda_k))_{1 \leq j, k \leq 18}.$$

Since Θ is a principal polarization on A , we have $Pf_{\Theta}(H_1(A, \mathbb{Z})) = 1$.

For $j \in \{0, \dots, 8\}$, it is easy to find a basis of the lattice Λ_j . We compute that

$$P_j^2 = a^{18} 19^{8-2j}$$

where P_j its Pfaffian of Λ_j . As a is positive, the only possibility that P_j equals 1 is $j = 4$ and $a = 1$. Moreover since all coefficients of the matrix M_{Λ_4} are integers, the alternating form $\frac{i}{\sqrt{19}} \sum dx_k \wedge d\bar{x}_k$ defines a principal polarization on A , see [6, Chap. 4.1]. $\square$

Theorem 1 follows from Lemmas 4 and 5.

Remark 6. One can compute that the involution μ acts on the principally polarized Abelian variety (A, Θ) , and therefore that $\text{Aut}(A, \Theta)$ contains the group $PSL_2(\mathbb{F}_{19})$.

Remark 7. In [5], Beauville proves that the intermediate Jacobian of the S_6 -symmetric quartic threefolds F is not a product of Jacobians of curves (see also [4]). That implies, using Clemens-Griffiths results for threefolds, the irrationality of F . Using the arguments at the end of [5], one can see that the principally polarized abelian ninefold (A, Θ) is not a product of Jacobians of curves.

2. INTERMEDIATE JACOBIANS OF THE ADLER CUBIC SEVENFOLD

2.1. Intermediate Jacobian. For a smooth algebraic complex manifold X of dimension $n = 2k + 1$, let $H^{p,q}(X)$ be the Hodge (p, q) -cohomology spaces $H^{p,q}(X) = H^q(X, \Omega^p)$, and $h^{p,q} = \dim H^{p,q}(X)$ be the Hodge numbers of X , $0 \leq p + q \leq 2n = 4k + 2$. The intersection of real $(2k + 1)$ -dimensional chains

in X defines an integer valued quadratic form Q_G on the integer cohomology $H_{2k+1}(X, \mathbb{Z})$, and yields an embedding of $H_{2k+1}(X, \mathbb{Z})$ as an integer lattice in the dual space of $H^{2k+1,0}(X) + H^{2k,1}(X) + \dots + H^{k+1,k}(X)$. The quotient compact complex torus

$$J_G(X) = (H^{2k+1,0}(X) + H^{2k,1}(X) + \dots + H^{k+1,k}(X))^* / H_{2k+1}(X, \mathbb{Z})$$

is the *Griffiths intermediate Jacobian* of the n -fold X , see [8, p.123]. By the Riemann-Hodge bilinear relations, the quadratic form Q_G is definite on any space $H^{p,n-p}(X)$, and has opposite signs on $H^{p,n-p}(X)$ and $H^{p',n-p'}(X)$ if and only if $p-p'$ odd. Moreover $\Im m(Q_G)$ takes integral values on $H_{2k+1}(X, \mathbb{Z})$. We call the pair $(J_G(X), Q_G)$ the polarized Griffiths intermediate Jacobian.

The quadratic form Q_G gives the polarized torus $(J_G(X), Q_G)$ a structure of a polarized Abelian variety if and only if p has the same parity for all non-zero spaces $H^{p,n-p}(X)$, ([8, p.123]).

2.2. Griffiths formulas. The middle Hodge groups of a smooth cubic sevenfold can be computed by the following Griffiths formulas, which we shall use below in order to determine the representation of the action of the symmetry group $PSL_2(\mathbb{F}_{19})$ on the middle cohomology of the Adler cubic sevenfold.

Let $X = (f(x) = 0)$ be a smooth hypersurface of degree $m \geq 2$ in the complex projective space $\mathbb{P}^{n+1}(x)$, $(x) = (x_1, \dots, x_{n+2})$. Let $S = \bigoplus_{d \geq 0} S_d$ be the graded polynomial ring $S = \mathbb{C}[x_0, \dots, x_{n+2}]$, with S_d being the space of polynomials of degree d . Let $I = \bigoplus I_d \subset S$ be the graded ideal generated by the $n+2$ partials $\frac{\partial f}{\partial x_j}$, $j = 1, \dots, n+2$, with $I_d = I \cap S_d$.

Let $R = S/I = \bigoplus R_d$ be the graded Jacobian ring of X (or the Jacobian ring of the polynomial $f(x)$), with graded components $R_d = S_d/I_d$. Then for $p+q=n$, the primitive cohomology group $H_{prim}^{p,q}(X)$ is isomorphic to the graded piece $R_{m(q+1)-n-2}$, where $R_d = 0$ for $d < 0$, ([8], p. 169). For odd $n = 2k+1$ all the middle cohomology of X are primitive, and in this case

$$H^{p,q}(X) \cong R_{m(q+1)-n-2}, \quad p+q = 2k+1 = \dim X, \quad m = \deg X.$$

In particular, for a smooth cubic sevenfold $X = (f(x) = 0) \subset \mathbb{P}^8$ one has

$$H^{7-q,q}(X) = R_{3(q+1)-9}, \quad 0 \leq q \leq 7,$$

which yields

$$\begin{aligned} H^{7,0}(X) &= R_{-6} = 0, \quad H^{6,1}(X) = R_{-3} = 0, \\ H^{5,2}(X) &= R_0 \cong \mathbb{C}, \quad H^{4,3}(X) = R_3 \cong \mathbb{C}^{84}. \end{aligned}$$

2.3. Character table of $PSL_2(\mathbb{F}_{19})$. In Proposition 8 below, we shall use the known description of the irreducible representations of the automorphism group $PSL_2(\mathbb{F}_{19}) \subset \text{Aut}(X)$, which we state here in brief:

By [3], the group $PSL_2(\mathbb{F}_{19})$ has 12 conjugacy classes:

$$1, \{w_1\}, \{w_2\}, \{x\}, \{x^2\}, \{x^3\}, \{x^4\}, \{y\}, \{y^2\}, \{y^3\}, \{y^4\}, \{y^5\}$$

where x has order 9, y has order 10, $w_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, $w_2 = w_1^2$ (we observe that $w_1^2, w_1^3 \in \{w_2\}$, $w_2^2, w_2^3 \in \{w_1\}$). Correspondingly, the 12 irreducible representations of $PSL_2(\mathbb{F}_{19})$ are

$$T_1, W_9, \overline{W}_9, W_{18}^1, W_{18}^2, W_{18}^3, W_{18}^4, W_{20}^1, W_{20}^2, W_{20}^3, W_{20}^4, W_{19}$$

where the notations are uniquely defined by the character table of $PSL_2(\mathbb{F}_{19})$ below for which $a_k = 2 \cos(\frac{2k\pi}{9})$, $b_k = -2 \cos(\frac{k\pi}{5})$ and $\nu = \frac{-1+i\sqrt{19}}{2}$. Note that the representation W_9 is described otherwise in section 1.

	1	w_1	w_2	x	x^2	x^3	x^4	y	y^2	y^3	y^4	y^5
1	1	1	1	1	1	1	1	1	1	1	1	1
W_9	9	ν	$\bar{\nu}$	0	0	0	0	1	-1	1	-1	1
$\overline{W}_9$	9	$\bar{\nu}$	ν	0	0	0	0	1	-1	1	-1	1
W_{18}^1	18	-1	-1	0	0	0	0	b_1	b_2	b_3	b_4	2
W_{18}^2	18	-1	-1	0	0	0	0	b_2	b_4	b_6	b_8	-2
W_{18}^3	18	-1	-1	0	0	0	0	b_3	b_6	b_9	b_{12}	2
W_{18}^4	18	-1	-1	0	0	0	0	b_4	b_8	b_{12}	b_{16}	-2
W_{20}^1	20	1	1	a_1	a_2	a_3	a_4	0	0	0	0	0
W_{20}^2	20	1	1	a_2	a_4	a_6	a_8	0	0	0	0	0
W_{20}^3	20	1	1	a_3	a_6	a_9	a_{12}	0	0	0	0	0
W_{20}^4	20	1	1	a_4	a_8	a_{12}	a_{16}	0	0	0	0	0
W_{19}	19	0	0	1	1	1	1	-1	-1	-1	-1	-1

FIGURE 2.1. Character table of $PSL_2(\mathbb{F}_{19})$

2.4. Periods of the Adler cubic. Let $X \subset \mathbb{P}^8 = \mathbb{P}(W_9)$ be the Adler cubic sevenfold

$$\{f_{19} = x_1^2 x_6 + x_6^2 x_2 + x_2^2 x_7 + x_7^2 x_4 + x_4^2 x_5 + x_5^2 x_8 + x_8^2 x_9 + x_9^2 x_3 + x_3^2 x_1 - 2(x_1 x_7 x_8 + x_2 x_3 x_5 + x_4 x_6 x_9) = 0\},$$

and let $H^{p,q} = H^{p,q}(X)$ be the Hodge cohomology spaces of X .

Proposition 8. *The representation of the group $PSL_2(\mathbb{F}_{19})$ on the 84-dimensional space $H^{4,3}(X)^*$ of the Adler cubic X is:*

$$H^{4,3}(X)^* = \overline{W}_9 \oplus W_{18}^1 \oplus W_{18}^3 \oplus W_{19} \oplus W_{20}^3.$$

The group $PSL_2(\mathbb{F}_{19})$ acts trivially on the one-dimensional space $H^{5,2}(X)^*$.

Remark 9. Let a group G acts on a vector space V on the left. Then the group G acts on the right on the dual space V^* : for $\ell \in V^*$ and $g \in G$, $\ell \cdot g = \ell \circ g$. Since the traces of the action of g on V and on V^* are equal, the two representations V and V^* are isomorphic. The representation V^* should not be confused with the dual representation of G defined such that the pairing between V and V^* is G -invariant (see [11]).

Proof. In order to decompose the representation of $PSL_2(\mathbb{F}_{19}) \subset Aut(X)$ on the dual cohomology space $H^{4,3*}$, we shall use the identification as representation space

$$H^{4,3}(X) \cong R_3 = S_3/I_3 = \text{Sym}^3 W_9/I_3$$

between $H^{4,3}(X)$ and the graded component of degree 3 in the quotient polynomial ring $S = \mathbb{C}[x_1, \dots, x_9]$ by the Jacobian ideal I spanned by the 9 partials of f_{19} , see above.

The space $\mathbb{P}^8$ containing the Adler cubic X is the projectivization of the representation space $W_9 = (S_1)^*$. We have $S_3 = \text{Sym}^3(W_9^*) \cong \text{Sym}^3(W_9)$. Let us decompose $\text{Sym}^3 W_9$ into irreducible representations of $PSL_2(\mathbb{F}_{19})$. The character of the third symmetric power of a representation V is:

$$\chi_{\text{Sym}^3 V}(g) = \frac{1}{6}(\chi_V(g)^3 + 3\chi_V(g^2)\chi_V(g) + 2\chi_V(g^3)),$$

see [16]. Therefore the traces of the action of the elements $1, w_1, w_2$ etc on $\text{Sym}^3 W_9$ are

$$v = {}^t(165, 3 - \nu, 3 - \bar{\nu}, 0, 0, 3, 0, 0, 0, 0, 5).$$

Using the character table of $PSL_2(\mathbb{F}_{19})$, we obtain:

$$\text{Sym}^3 W_9 = T_1 \oplus \overline{W}_9 \oplus W_{18}^1 \oplus W_{18}^3 \oplus W_{20}^1 \oplus W_{20}^2 \oplus (W_{20}^3)^{\oplus 2} \oplus W_{20}^4 \oplus W_{19}.$$

The graded component I_2 of the Jacobian ideal I of X is generated by the 9 derivatives $\frac{df_{19}}{dx_k}$, $k = 1, \dots, 9$. The space I_2 is a representation of $PSL_2(\mathbb{F}_{19})$. The action of w_1 on x_j is the multiplication by ξ^{j^2} , where $\xi = e^{2i\pi/19}$, see section 1. The action of w_1 on I_2 is then easy to compute. By example, since

$$\frac{df_{19}}{dx_1} = 2x_1x_6 + x_3^2 - 2x_7x_8,$$

we get $w_1 \cdot \frac{df_{19}}{dx_1} = \xi^{-1} \frac{df_{19}}{dx_1}$. By looking at the character table, we obtain that the representation I_2 is $\overline{W}_9$. Therefore $I_3 = W_9 \otimes \overline{W}_9$. Using the fact that for two representations V_1, V_2 , their characters satisfy the relation $\chi_{V_1 \otimes V_2} = \chi_{V_1} \chi_{V_2}$, we obtain:

$$I_3 = W_9 \otimes \overline{W}_9 = T_1 \oplus W_{20}^1 \oplus W_{20}^2 \oplus W_{20}^3 \oplus W_{20}^4.$$

Since

$$H^{4,3}(X)^* \cong (S_3/I_3)^* \cong \text{Sym}^3 W_9/I_3$$

then

$$H^{4,3}(X)^* = \overline{W}_9 \oplus W_{18}^1 \oplus W_{18}^3 \oplus W_{19} \oplus W_{20}^3.$$

The action of the group $PSL_2(\mathbb{F}_{19}) \subset Aut(X)$ on $H^{5,2}(X)^* \cong R_0^* \cong \mathbb{C}$ is trivial, see the character table. $\square$

Let V be a representation of a finite group G and let W be an irreducible representation of G of character χ_W . We know that there exist a uniquely determined integer $a \geq 0$ and a representation V' of G such that W is not a

sub-representation of V' and V is (isomorphic to) $W^{\oplus a} \oplus V'$. By [11] formula (2.31) p. 23, the linear endomorphism

$$\psi_W = \frac{\dim W}{|G|} \sum_{g \in G} \overline{\chi_W(g)} g : V \rightarrow V$$

is the projection of V onto the factor consisting of the sum of all copies of W appearing in V i.e. is the projection onto $W^{\oplus a}$. For the trivial representation T , $\chi_T = 1$ and ψ_T is the projection of V onto the invariant space V^G .

Recall that an endomorphism h of a torus Y acts on the tangent space TY of Y by a linear endomorphism dh called the analytic representation of h . The kernel of the map $End(Y) \rightarrow End(TY)$, $h \rightarrow dh$ is the group of translations (see [6]) ; in the following we will work with endomorphisms up to translation.

Suppose that for some irreducible representations $W_1, \dots, W_k$ the sum

$$\overline{\chi_{W_1}(g)} + \dots + \overline{\chi_{W_k}(g)}$$

is an integer for every g . Then the endomorphism

$$h = \sum_{g \in G} (\overline{\chi_{W_1}(g)} + \dots + \overline{\chi_{W_k}(g)}) g \in End(J_G X)$$

is well defined and its analytic representation is

$$dh = \sum_{g \in G} (\overline{\chi_{W_1}(g)} + \dots + \overline{\chi_{W_k}(g)}) dg \in End(TJ_G X).$$

The tangent space of the image of h (translated to 0) is the image of dh .

Corollary 10. *The torus $J_G X$ has the structure of an Abelian variety and is isogenous to:*

$$E \times A_9 \times A_{36} \times A_{19} \times A_{20}$$

where A_k is a k -dimensional Abelian subvariety of $J_G X$ and $E \subset J_G X$ an elliptic curve. The group $PSL_2(\mathbb{F}_{19})$ acts nontrivially on each factors $A_9, A_{36}, A_{19}, A_{20}$.

Proof. The character of the trivial representation is $\chi_0 = 1$. Let $\chi_1, \chi_2, \chi_3, \chi_4$ be respectively the characters of the representations $W_9 \oplus \overline{W}_9$, $W_{18}^1 \oplus W_{18}^2 \oplus W_{18}^3 \oplus W_{18}^4$, W_{19} and $W_{20}^1 \oplus W_{20}^2 \oplus W_{20}^3 \oplus W_{20}^4$. By the character table, the numbers $\chi_k(g)$, $g \in PSL_2(\mathbb{F}_{19})$ are integers, therefore we can define

$$q_k = \sum_{g \in PSL_2(\mathbb{F}_{19})} \chi_k(g) g$$

for $k = 0, \dots, 4$. The analytic representation of q_k is a multiple of the projection onto the subspace $(H^{5,2})^*, W_9, \dots$ of $TJ_G X$. The images of q_k , $k = 0, \dots, 4$ are therefore respectively 1, 9, 36, 19, 20-dimensional sub-tori of $J_G X$, stable by the action of the group ring $\mathbb{Z}[PSL_2(\mathbb{F}_{19})]$ and denoted respectively by $E, A_9, \dots, A_{20}$.

The image of the endomorphism $q_1 + \dots + q_4$ (resp. q_0) is a sub-torus whose tangent space is $H^{4,3}(X)^*$ (resp. $H^{5,2}(X)^*$), therefore $H_7(X, \mathbb{Z}) \cap H^{4,3}(X)^*$ is a lattice in $H^{4,3}(X)^*$ (resp. $H_7(X, \mathbb{Z}) \cap H^{5,2}(X)^*$ is a lattice in $H^{5,2}(X)^*$).

Let Q_G be the Hodge-Riemann form on the tangent space $(H^{5,2} \oplus H^{4,3})^*$. It is positive definite on $H^{5,2}(X)^*$, negative definite on $H^{4,3}(X)^*$, the space $H^{4,3}(X)^*$ is orthogonal to $H^{5,2}(X)^*$ with respect to Q_G and Q_G takes integral values on $H_7(X, \mathbb{Z}) \subset (H^{5,2} \oplus H^{4,3})^*$ (see [8], 114).

Let us define the quadratic form Q' on $(H^{5,2} \oplus H^{4,3})^*$ by $Q' = -Q_G$ on $H^{4,3}(X)^*$ and $Q' = Q_G$ on $H^{5,2}(X)^*$. This Q' is a definite quadratic form that takes integral values on the lattice $\Lambda \subset H^{5,2}(X)^*$ generated by $H_7(X, \mathbb{Z}) \cap H^{4,3}(X)^*$ and $H_7(X, \mathbb{Z}) \cap H^{5,2}(X)^*$. We thus see that $J' = (H^{5,2} \oplus H^{4,3})^* / \Lambda$ is an Abelian variety, and since $J_G X$ is isogenous to J' , $J_G X$ is also an Abelian variety. $\square$

3. ON THE ADLER-KLEIN PENCIL OF CUBICS.

Here we study the Adler-Klein pencil of cubics $X_\lambda = \{f_\lambda = 0\}$,

$$f_\lambda = x_1^2 x_6 + x_6^2 x_2 + x_2^2 x_7 + x_7^2 x_4 + x_4^2 x_5 + x_5^2 x_8 + x_8^2 x_9 + x_9^2 x_3 + x_3^2 x_1 \\ + \lambda(x_1 x_7 x_8 + x_2 x_3 x_5 + x_4 x_6 x_9),$$

where X_{-2} is the Adler cubic, X_0 is the Klein cubic. Since X_{-2} and X_0 are smooth, the general member of the pencil is smooth.

The automorphism group of X_λ contains the group

$$H = \mathbb{Z}/9\mathbb{Z} \rtimes \mathbb{Z}/19\mathbb{Z}$$

whose law is defined multiplicatively by:

$$(a, b)(c, d) = (a + c, 4^c \cdot b + d).$$

Remark. The group H is a subgroup of $PSL_2(\mathbb{F}_{19})$: it is the stabilizer of a point in the projective line $\mathbb{P}^1(\mathbb{F}_{19})$ for the action of the simple group $PSL_2(\mathbb{F}_{19})$. Since $\mathbb{P}^1(\mathbb{F}_{19})$ has 20 points, there are therefore 20 such subgroups in $PSL_2(\mathbb{F}_{19})$.

Let us study the representation of H on the tangent space of the intermediate Jacobian of X_λ . Let be $a = (1, 0)$ and $b = (0, 1) \in H$. Let us denote by C_g the conjugacy class of an element $g \in H$. The conjugacy classes of the group H are the 11 classes: C_b , C_{b^2} and C_{a^k} , $k = 0, \dots, 8$. Let μ be a 9th-primitive root of unity. The group $\mathbb{Z}/9\mathbb{Z}$ is the quotient of H by $\mathbb{Z}/19\mathbb{Z}$, thus the irreducible one-dimensional representation

$$\chi_k : \begin{cases} \mathbb{Z}/9\mathbb{Z} & \rightarrow \mathbb{C}^* \\ a & \rightarrow \mu^{ka} \end{cases}, k \in \{0, \dots, 8\}$$

induces an irreducible representation V_k of H . Let $V_9, \overline{V}_9$ be the restrictions of the representations $W_9, \overline{W}_9$ to $H \subset PSL_2(\mathbb{F}_{19})$. As $171 = 1^2 + \dots + 1^2 + 9^2 + 9^2$ is equal to the order of H , the representations $V_0, \dots, V_8, V_9, \overline{V}_9$ are the 11 irreducible non-isomorphic representations of H (see [11]).

Proposition 11. *The restrictions of the representations W_{18}^3, W_{19}, W_{20} of $PSL_2(\mathbb{F}_{19})$ to $H \subset PSL_2(\mathbb{F}_{19})$ are decomposed as follows:*

$$\begin{aligned} W_{18}^3 &= V_9 + \overline{V}_9 \\ W_{19} &= V_0 + V_9 + \overline{V}_9 \\ W_{20} &= V_3 + V_6 + V_9 + \overline{V}_9. \end{aligned}$$

	1	a	a^2	a^3	a^4	a^5	a^6	a^7	a^8	b	b^2
V_0	1	1	1	1	1	1	1	1	1	1	1
V_1	1	μ	μ^2	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	μ^8	1	1
V_2	1	μ^2	μ^4	$\dots$					$\vdots$	1	1
V_3	1	$\vdots$		$\ddots$						1	1
V_4	1	$\vdots$								1	1
V_5	1	$\vdots$								1	1
V_6	1	$\vdots$								1	1
V_7	1	$\vdots$	$\vdots$					$\ddots$	$\vdots$	1	1
V_8	1	μ^8	μ^{16}	$\dots$	$\dots$	$\dots$	$\dots$	$\dots$	μ^{64}	1	1
V_9	9	0	0	0	0	0	0	0	0	ν	$\bar{\nu}$
$\overline{V}_9$	9	0	0	0	0	0	0	0	0	$\bar{\nu}$	ν

FIGURE 3.1. Character table of $H = \mathbb{Z}/9\mathbb{Z} \rtimes \mathbb{Z}/19\mathbb{Z}$. $\nu = \frac{1}{2}(-1 + i\sqrt{19})$, $\mu^9 = 1$, $a = (1, 0)$, $b = (0, 1)$.

Proof. There exist positive integers $a_0, \dots, a_9, b_0$ such that the representation of H on W_{18}^3 equals (in fact, is isomorphic) to $V_0^{\oplus a_0} \oplus \dots \oplus V_9^{\oplus a_9} \oplus \overline{V}_9^{\oplus b_9}$. For an element $g \in H$ acting on a vector space V , we denote by $Tr(g|V)$ the trace of g acting on V . We have

$$Tr(g|W_{18}^3) = a_0 Tr(g|V_0) + \dots + a_9 Tr(g|V_9) + b_9 Tr(g|\overline{V}_9).$$

The values of $Tr(g|V_1), \dots, Tr(g|V_9), Tr(g|\overline{V}_9)$ are given in the above character table, and are independent of the conjugacy class of g . Since we know the values of $Tr(g|W_{18}^3)$, $g \in H$ from the character table of $PSL_2(\mathbb{F}_{19})$, by elementary linear algebra we obtain that $a_0 = \dots = a_8 = 0$ and $a_9 = b_9 = 1$, therefore $W_{18}^3 = V_9 \oplus \overline{V}_9$. In the same way, we find $W_{19} = V_0 + V_9 + \overline{V}_9$ and $W_{20} = V_3 + V_6 + V_9 + \overline{V}_9$. $\square$

Let X_λ be a smooth cubic in the Klein-Adler pencil and let $J_G X_\lambda$ be the Griffiths intermediate Jacobian of X_λ .

Corollary 12. *The representation of H on the tangent space to the Griffiths intermediate Jacobian of X_λ is:*

$$TJ_G X_\lambda = V_0 + H^{4,3}(X)^* = 2V_0 + V_3 + V_6 + 4V_9 + 5\overline{V}_9.$$

There exist subtori A_2, B_2, B_{81} of $J_G X_\lambda$ of dimension respectively 2, 2, 81, such that B_2 and B_{81} are Abelian varieties and such that there is an isogeny of complex tori

$$J_G X_\lambda \rightarrow A_2 \times B_2 \times B_{81}.$$

Proof. For the Adler cubic X_{Ad} , the representation of $PSL_2(\mathbb{F}_{19})$ on $H^{4,3}(X_{Ad})$ is

$$H^{4,3}(X_{Ad}) = \overline{W}_9 \oplus W_{18}^1 \oplus W_{18}^3 \oplus W_{19} \oplus W_{20}^3.$$

From Proposition 11, we know the representation of H on each of the factors of $H^{4,3}$. Since H acts also on each X_λ , the representation of H on $H^{4,3}(X_\lambda)$ is the same for all cubics X_λ in the pencil, and we have:

$$R_3 = S^3 V_9 / I_3 = V_0 + V_3 + V_6 + 4V_9 + 5\overline{V}_9.$$

Thus $TJ_G X_\lambda = (V_0)^{\oplus 2} \oplus V_{83}$. The image of the endomorphism $\sum_{g \in G} g$ is therefore a 2-dimensional torus A_2 . By considering the quotient map $J_G X_\lambda \rightarrow J_G X_\lambda / A_2$, we see that $V_{83} \cap H_7(X_\lambda, \mathbb{Z})$ is a lattice. There is moreover on it a positive definite integral valued form (the restriction of the Hodge-Riemann form), therefore V_{83} is the tangent space of a 83-dimensional Abelian subvariety A_{83} of $J_G X_\lambda$. By using the same arguments as in section 2, we see that A_{83} is isogenous to a product $B_2 \times B_{81}$ of two abelian sub-varieties. $\square$

REFERENCES

- [1] Adler A., *On the automorphism group of certain hypersurfaces*, J. Algebra **72** (1981), no.1, 146–165.
- [2] Adler A., *Cubic Invariants for $SL_2(\mathbb{F}_q)$* , J. Algebra **144** (1992), 178–186.
- [3] Conway, J. H., Curtis, R. T., Norton, S. P., Parker, R. A., Wilson, R. A., *Atlas of finite groups*, Oxford University Press, 1985, xxxiv+252 pp
- [4] Beauville A., *Non-rationality of the symmetric sextic Fano threefold*, Geometry and Arithmetic, EMS Congress Reports, (2012), pp. 57–60.
- [5] Beauville A., *Non-rationality of the S_6 -symmetric quartic threefolds*, arXiv 1212.5345
- [6] Birkenhake C., Lange H., *Complex abelian varieties*, Grundlehren, Vol. **302**, 2nde edition, Springer 2004.
- [7] Albano A., Collino A., *On the Griffiths group of the cubic sevenfold*, Math. Ann. **299**, 715–726.
- [8] Carlson J., Green M., Griffiths P., Harris J., *Infinitesimal variation of Hodge structure I*, Compositio Math. **50** (1983), no. 2-3, 109–205.
- [9] Candelas P., de la Ossa X., van Geemen B., van Straten D., *Lines on the Dwork Pencil of Quintic Threefolds*, arXiv 1206.4961
- [10] Donagi R., *Generic Torelli for projective hypersurfaces*, Compositio Math. **50** (1983), no. 2-3, 325–353.
- [11] Fulton W., Harris J., *Representation theory, A first course*, GTM **129**, Springer-Verlag, New York, 1991.
- [12] Griffiths, P.A. *On the periods of certain rational integrals I, II*, Ann. of Math. (2) **90** (1969), 460–495; *ibid.* (2) **90** (1969) 496–541.
- [13] Iliev A., Manivel L., *Fano manifolds of Calabi-Yau type*, arXiv :1102.3623
- [14] Klein F., *Über die Transformation elfter Ordnung der elliptischen Functionen*, Math. Annalen, vol. **15**, (1879) no. 3, pp. 533–555.

- [15] Masley J., *Solution of the class number two problem for cyclotomic fields*, Invent. Math. **28** (1975), 243–244.
- [16] Pulay P., Zhou X.-F., (1989), *Characters for symmetric and antisymmetric higher powers of representations: Application to the number of anharmonic force constants in symmetrical molecules*, J. Comput. Chem. **10**, 935–938.
- [17] Roulleau X., *The Fano surface of the Klein cubic threefold*, J. Math. Kyoto Univ. **49** (2009), no. 1, 113–129.

Atanas Iliev
Department of Mathematics
Seoul National University
151-747 Seoul, Korea
e-mail: ailiev@snu.ac.kr

Xavier Roulleau
Laboratoire de Mathématiques et Applications,
Téléport 2 - BP 30179 -
Boulevard Pierre et Marie Curie
86962 Futuroscope Chasseneuil
France
e-mail: xavier.roulleau@math.univ-poitiers.fr